

Absolute values, valuations and completion

F. Crivelli (flcrivel@student.ethz.ch)

April 21, 2008

Introduction

During this talk I'll introduce the basic definitions and some results about valuations, absolute values of fields and completions. These notions will give two basic examples: the *rational function field* $\mathbb{F}_q(T)$ and the field \mathbb{Q}_p of the *p-adic numbers*.

1 Absolute values and valuations

All along this section, K denote a field.

1.1 Absolute values

We begin with a well-known

Definition 1. An *absolute value* of K is a function

$$|\cdot| : K \rightarrow \mathbb{R}$$

satisfying these properties, $\forall x, y \in K$:

1. $|x| = 0 \Leftrightarrow x = 0$,
2. $|x| \geq 0$,
3. $|xy| = |x||y|$,
4. $|x + y| \leq |x| + |y|$. (Triangle inequality)

Note that if we set $|x| = 1$ for all $0 \neq x \in K$ and $|0| = 0$, we have an absolute value on K , called the *trivial absolute value*. From now on, when we speak about an absolute value $|\cdot|$, we assume that $|\cdot|$ is non-trivial. Moreover, if we define $d : K \times K \rightarrow \mathbb{R}$ by $d(x, y) = |x - y|$, $x, y \in K$, d is a metric on K and we have a topological structure on K . We have also some basic properties that we can deduce directly from the definition of an absolute value.

Lemma 1. Let $|\cdot|$ be an absolute value on K . We have

1. $|1| = 1$,
2. $|\zeta| = 1$, for all $\zeta \in K$ with $\zeta^d = 1$ for some $0 \neq d \in \mathbb{N}$ (ζ is a root of unity),
3. $|x^{-1}| = |x|^{-1}$,
4. $||x| - |y|| \leq |x - y|$.

Proof:

1. $|1|^2 = |1^2| = |1| \Rightarrow |1| = 1$.
2. $|\zeta|^d = |\zeta^d| = |1| = 1 \Rightarrow |\zeta| = 1$.
3. $1 = |xx^{-1}| = |x||x^{-1}| \Rightarrow |x^{-1}| = |x|^{-1}$,
4. Set $a := x - y$, $b := y$. We have $|a + b| \leq |a| + |b|$, therefore $|x| - |y| = |a + b| - |b| \leq |a| = |x - y|$.

□

Another definition about absolute values.

Definition 2. Two absolute values on K are **equivalent** if they define the same topology on K .

The next theorem and its corollary give us other conditions to verify if two absolute values are equivalent.

Theorem 1. Let $|\cdot|_1$ and $|\cdot|_2$ be two absolute values on K . They are equivalent if, and only if, there exists $s \geq 0$ real such that

$$|x|_1 = |x|_2^s, \forall x \in K.$$

Proof: If we have $|x|_1 = |x|_2^s$, $s > 0$, clearly the two absolute values are equivalent since they define the same open sets.

If $|\cdot|_1$ and $|\cdot|_2$ are equivalent, a series converging to 0 with respect to $|\cdot|_1$ will converge to 0 with respect to $|\cdot|_2$. Moreover, for all K field and $|\cdot|$ on K , the inequality $|x| < 1$ is equivalent to saying that the sequence $\{x^n\}_{n \in \mathbb{N}}$ converges to 0. Therefore if $|\cdot|_1$ and $|\cdot|_2$ are equivalent, we have

$$|x|_1 < 1 \Leftrightarrow |x|_2 < 1.$$

Let $y \in K$ be an element such that $|y|_1 > 1$ and let $x \in K$, $x \neq 0$. Then there exists $\alpha \in \mathbb{R}$ such that $|x|_1 = |y|_1^\alpha$. Let $\left\{ \frac{m_i}{n_i} \right\}_{i \in \mathbb{N}}$ be a sequence of rational numbers

($m_i \in \mathbb{Z}$, $n_i \in \mathbb{N}^*$) converging to α from above. Then $|x|_1 = |y|_1^\alpha < |y|_1^{\frac{m_i}{n_i}}$ and hence

$$\left| \frac{x^{n_i}}{y^{m_i}} \right|_1 < 1 \Rightarrow \left| \frac{x^{n_i}}{y^{m_i}} \right|_2 < 1.$$

This gives $|x|_2 < |y|_2^{\frac{m_i}{n_i}}$ and consequently $|x|_2 \leq |y|_2^\alpha$. Taking a sequence $\left\{ \frac{m_i}{n_i} \right\}_{i \in \mathbb{N}}$ converging to α from below will give $|x|_2 \geq |y|_2^\alpha$ and therefore $|x|_2 = |y|_2^\alpha$. So, for all $0 \neq x \in K$, we have

$$\frac{\log |x|_1}{\log |x|_2} = \frac{\log |y|_1}{\log |y|_2} =: s$$

and hence $|x|_1 = |x|_2^s$. Finally, $|y|_1 > 1$ implies $|y|_2 > 1$ and so $s > 0$.

□

Corollary. *Two absolute values $| \cdot |_1$ and $| \cdot |_2$ on K are equivalent if, and only if,*

$$|x|_1 < 1 \iff |x|_2 < 1.$$

We continue with this important

Definition 3. *An absolute value is called **non-Archimedean** if we have*

$$|x + y| \leq \max\{|x|, |y|\}, \forall x, y \in K.$$

*Otherwise the absolute value is called **Archimedean**.*

Note that if $| \cdot |$ is non-Archimedean, for $x, y \in K$ we have

$$|x| \neq |y| \implies |x + y| = \max\{|x|, |y|\}.$$

Indeed: w.l.o.g we can assume $|x| > |y|$, then obviously $|x + y| \leq \max\{|x|, |y|\} = |x|$.

On the other side, $|x| = |x - y + y| \leq \max\{|x - y|, |y|\}$. Assume that $\max\{|x + y|, |y|\} = |y|$, then $|x| \leq |y| < |x|$. This contradicts the hypothesis $|x| > |y|$, hence $\max\{|x + y|, |y|\} = |x + y|$ and so $|x| \leq |x + y|$.

1.2 Valuations

We introduce the symbol ∞ with the convention that for all $a \in \mathbb{R}$ we have $a < \infty$, $a + \infty = \infty$ and $\infty + \infty = \infty$. As for absolute values, we start with a basic

Definition 4. *A **valuation** on K is a function*

$$v : K \rightarrow \mathbb{R} \cup \{\infty\}$$

satisfying these properties, for all $x, y \in K$:

1. $v(x) = \infty \Leftrightarrow x = 0$,
2. $v(xy) = v(x) + v(y)$,
3. $v(x + y) \geq \min\{v(x), v(y)\}$.

Note that if we set $v(x) = 0$ for all $0 \neq x \in K$ and $v(0) = \infty$, we have a valuation on K , called the *trivial valuation*. From now on, when we speak about a valuation v , we assume that v is non-trivial.

Now, a lemma with some basic properties induced by the definition of valuation

Lemma 2. *Let v be a valuation on K . We have*

1. $v(1) = 0$,
2. $v(\zeta) = 0$, for all $\zeta \in K$ root of unity,
3. $v(x^{-1}) = -v(x), \forall x \in K^*$,
4. if $x, y \in K$ and $v(x) \neq v(y)$, $v(x + y) = \min\{v(x), v(y)\}$.

Proof:

1. $v(1) = v(1^2) = v(1) + v(1) \Rightarrow v(1) = 0$.
2. Let $\zeta \in K$ with $\zeta^d = 1$ for some $0 \neq d \in \mathbb{N}$. We have $dv(\zeta) = v(\zeta^d) = v(1) = 0$, therefore $v(\zeta) = 0$.
3. $0 = v(1) = v(xx^{-1}) = v(x) + v(x^{-1}) \Rightarrow v(x^{-1}) = -v(x)$.
4. W.l.o.g. we can assume that $v(x) > v(y)$ and so $y \neq 0$. If $x = 0$, obvious. Assume $x \neq 0$. Then $v(x + y) \geq \min\{v(x), v(y)\} = v(y)$. We have also $v(y) = v(y + x - x) \geq \min\{v(x + y), v(x)\}$. Assume that $\min\{v(x + y), v(x)\} = v(x)$, then $v(x) > v(y) \geq v(x)$ and we get a contradiction. So, $\min\{v(x + y), v(x)\} = v(x + y)$ and finally $v(y) \geq v(x + y)$.

□

Some further terminology regarding valuations.

Definition 5. *A valuation v on K is called **discrete** if $v(K^*) = s\mathbb{Z}$, for a real $s > 0$. Moreover, v is **normalized** if $s = 1$.*

We introduce now the equivalence between valuations.

Definition 6. *Two valuations v_1 and v_2 on K are **equivalent** if there exists a real $s > 0$ such that $v_1 = sv_2$.*

Note that if we have a discrete valuation on K with $v(K^*) = s\mathbb{Z}$, dividing it by s we obtain an equivalent normalized valuation.

1.3 Relations between non-Archimedean absolute values and valuations

The following theorem provides a relation between the non-Archimedean absolute values and the valuations on K .

Theorem 2. *Let $|\cdot|$ be a non-Archimedean absolute value on K and $s \in \mathbb{R}$, $s > 0$, then the function*

$$v_s : K \longrightarrow \mathbb{R} \cup \{\infty\}$$

$$x \longmapsto \begin{cases} -s \log |x| & \text{if } x \neq 0 \\ \infty & \text{if } x = 0 \end{cases}$$

is a valuation on K . Furthermore, if $s, s' \in \mathbb{R}$, $s, s' > 0$ and $s \neq s'$, v_s is equivalent to $v_{s'}$. Conversely, if v is a valuation on K and $q \in \mathbb{R}$, $q > 1$, the function

$$|\cdot|_q : K \longrightarrow \mathbb{R}$$

$$x \longmapsto \begin{cases} q^{-v(x)} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$$

is an absolute value on K . Besides, if $q, q' \in \mathbb{R}$, $q, q' > 1$ and $q \neq q'$, $|\cdot|_q$ is equivalent to $|\cdot|_{q'}$.

Proof: We just need to check the definitions of an absolute value and a valuation. We start with v_s . Clearly we have that $v_s(x) = \infty$ if, and only if, $x = 0$. Let $x, y \in K$, if $x = 0$ or $y = 0$, $xy = 0$ and $\infty = v_s(xy) = v(x) + v(y) = \infty$. Assume $x, y \neq 0$, then

$$v_s(xy) = -s \log |xy| = -s \log (|x| |y|) = -s \log |x| - s \log |y| = v_s(x) + v_s(y).$$

Let again be $x, y \in K$. If $x = y = 0$, then $\infty = v_s(x + y) = \min \{v_s(x), v_s(y)\} = \infty$. If $x = 0$, $y \neq 0$ (or $y = 0$, $x \neq 0$), $v_s(x + y) = v_s(y) = \min \{v_s(x), v_s(y)\}$ (or $v_s(x + y) = v_s(x) = \min \{v_s(x), v_s(y)\}$). Assume now $x, y \neq 0$. We have

$$\begin{aligned} v_s(x + y) &= -s \log |x + y| \\ &\geq -s \log (\max \{|x|, |y|\}) \\ &= \min \{-s \log |x|, -s \log |y|\} \\ &= \min \{v_s(x), v_s(y)\}. \end{aligned}$$

Therefore v_s is a valuation on K . Assume now $s, s' > 0$, $s \neq s'$. For all $0 \neq x \in K$, we have

$$v_s(x) = -s \log |x| = \left(\frac{s}{s'}\right) (-s' \log |x|) = \frac{s}{s'} v_{s'}(x).$$

This means that v_s and $v_{s'}$ are equivalent. We continue with $|\cdot|_q$. Clearly we have that $|x|_q = 0$ if, and only if, $x = 0$ and since $q > 1 > 0$ that $|x|_q \geq 0$ for all $x \in K$. Let $x, y \in K$, if $x = 0$ or $y = 0$, $xy = 0$ and $0 = |xy|_q = |x|_q |y|_q = 0$. Assume $x, y \neq 0$, then

$$|xy|_q = q^{-v(xy)} = q^{-v(x)-v(y)} = q^{-v(x)} q^{-v(y)} = |x|_q |y|_q.$$

Since $\max\{|x|_q, |y|_q\} \leq |x|_q + |y|_q$, it suffices to show that $|x + y|_q \leq \max\{|x|_q, |y|_q\}$. Let again be $x, y \in K$. If $x = y = 0$, $0 = |x + y|_q = \max\{|x|_q, |y|_q\} = 0$. If $x = 0, y \neq 0$ (or $y = 0, x \neq 0$), $|x + y|_q = |y|_q = \max\{|x|_q, |y|_q\}$ (or $|x + y|_q = |x|_q = \max\{|x|_q, |y|_q\}$). Assume now $x, y \neq 0$. We have

$$\begin{aligned} |x + y|_q &= q^{-v(x+y)} \\ &\leq q^{-\min\{v(x), v(y)\}} \\ &= \max\{q^{-v(x)}, q^{-v(y)}\} \\ &= \max\{|x|_q, |y|_q\}. \end{aligned}$$

Therefore, $|\cdot|_q$ is a non-Archimedean absolute value on K . Assume now $q, q' > 1$, $q \neq q'$ and set $r := \frac{\log q}{\log q'}$. For all $0 \neq x \in K$, we have

$$|x|_q = q^{-v(x)} = q'^{-rv(x)} = |x|_{q'}^r.$$

Consequently, $|\cdot|_q$ and $|\cdot|_{q'}$ are equivalent. □

From now on, when we will deal with fields with a non-Archimedean absolute value, according to theorem 2, we will just speak of a field with a valuation.

Remark on terminology: Note that some authors use the term “exponential valuation” rather than “valuation”. In this case the term “valuation” means “absolute value”.

1.4 Valuation ring and residue field

Theorem 3. *Let K be a field, v be a valuation on K and denote by $|\cdot|$ a corresponding non-Archimedean absolute value. Then:*

1. *the set*

$$\mathcal{O} := \{x \in K \mid v(x) \geq 0\} = \{x \in K \mid |x| \leq 1\}$$

*is an integral domain and a maximal proper subring of K , called the **valuation ring**; moreover, for all $0 \neq x \in K$, we have that $x \in \mathcal{O}$ or $x^{-1} \in \mathcal{O}$,*

2. *the set*

$$\mathcal{O}^* := \{x \in K \mid v(x) = 0\} = \{x \in K \mid |x| = 1\}$$

is the group of units of \mathcal{O} ,

3. the set

$$\mathfrak{p} := \mathcal{O} \setminus \mathcal{O}^* = \{x \in K \mid v(x) > 0\} = \{x \in K \mid |x| < 1\} = \{x \in \mathcal{O} \mid x^{-1} \notin \mathcal{O}\}$$

is the unique maximal ideal of \mathcal{O} .

A ring that has a unique maximal ideal is called a **local ring**. By 3., \mathcal{O} is a local ring. Besides, two equivalent valuations (or two equivalent non-Archimedean absolute values) on K give the same valuation ring.

Proof: We just consider the valuation v on K , since by theorem 2 the valuations and the non-Archimedean absolute values are closely related.

It is easy to check that if two valuation v and v' on K are equivalent, i.e., there is a positive real s such that $v = sv'$, the sets \mathcal{O} , \mathcal{O}^* and \mathfrak{p} are the same since if $v(x) \geq 0$, resp. $v(x) = 0$, resp. $v(x) > 0$, $v'(x) = s^{-1}v(x) \geq 0$, resp. $v'(x) = s^{-1}v(x) = 0$, resp. $v'(x) = s^{-1}v(x) > 0$.

To prove that \mathcal{O} is a integral domain, it suffices to show that is closed under addition and multiplication and that every element in \mathcal{O} has an additive inverse in \mathcal{O} , since all the remaining properties are verified on the field K and therefore also for the subset \mathcal{O} . Clearly, $0, 1 \in \mathcal{O}$ since $v(0) = \infty > v(1) = 0 \geq 0$. Take $0 \neq x \in \mathcal{O}$ and so $-x \in K$. We have $\infty = v(0) = v(x - x) > \min\{v(x), v(-x)\}$ since $x \neq 0$ (and therefore $-x \neq 0$). By property 3. of Lemma 2, this implies that $v(-x) = v(x) \geq 0$, hence $-x \in \mathcal{O}$. Let $x, y \in \mathcal{O}$. Then $v(xy) = v(x) + v(y) \geq 0$ and so $xy \in \mathcal{O}$.

Similarly, $v(x + y) \geq \min\{v(x), v(y)\} \geq 0$ and so $x + y \in \mathcal{O}$. Take now $0 \neq x \in K$. If $v(x) \geq 0$, $x \in \mathcal{O}$, and if $v(x) < 0$, $v(x^{-1}) = -v(x) > 0$ and hence $x^{-1} \in \mathcal{O}$. Let now be $z \in K \setminus \mathcal{O}$. We want to show that $\mathcal{O}[z] = K$ in order to prove that \mathcal{O} is a maximal proper subring of K . Since $z \notin \mathcal{O}$, $z^{-1} \in \mathcal{O}$ and $v(z^{-1}) > 0$. Take an element $y \in K$, then there exists $k \in \mathbb{N}$ such that $v(yz^{-k}) \geq 0$ since $v(z^{-1}) > 0$. Consequently, $w := yz^{-k} \in \mathcal{O}$ and finally $y = wz^k \in \mathcal{O}[z]$ and so $\mathcal{O}[z] = K$.

Assume now $x \in \mathcal{O}$ and $x^{-1} \in \mathcal{O}$. If $v(x) > 0$, $v(x^{-1}) = -v(x) < 0$, therefore $v(x) = 0$. The other inclusion is obvious. This shows that \mathcal{O}^* is the group of units of \mathcal{O} .

We show now that \mathfrak{p} is the unique maximal ideal of \mathcal{O} . Let $x \in \mathfrak{p}$ and $z \in \mathcal{O}$. Then $v(zx) = v(z) + v(x) \geq v(x) > 0$ and hence $zx \in \mathfrak{p}$. Let now $x, y \in \mathfrak{p}$, then $v(x - y) \geq \min\{v(x), v(-y)\}$. If $y = 0$, $x - y \in \mathfrak{p}$. If $y \neq 0$, we have that $\infty = v(0) = v(y - y) > \min\{v(y), v(-y)\}$ and, as before, $v(y) = v(-y) > 0$. Therefore $v(x - y) > 0$. This means that $x - y \in \mathfrak{p}$ and hence \mathfrak{p} is an additive subgroup of \mathcal{O} . This shows that \mathfrak{p} is an ideal of \mathcal{O} . Assume now there exists an ideal A of \mathcal{O} such that $\mathfrak{p} \subsetneq A$. Then there exists $x \in \mathcal{O}^* \cap A$. Therefore $1 \in A$ and $A = \mathcal{O}$. This shows that \mathfrak{p} is maximal. Assume now that there is a maximal ideal $B \neq \mathcal{O}$ of \mathcal{O} such that $B \neq \mathfrak{p}$. We must have that $B \cap \mathcal{O}^* = \{0\}$, because if not, $1 \in B$ and so $B = \mathcal{O}$. This implies that $B \subsetneq \mathfrak{p}$ and therefore B is not maximal.

□

The theorem tells us that for two equivalent valuation we have the same valuation ring, so, from now on, if a valuation is discrete, we can assume without loss of generality that it is normalized. Clearly if a result holds for a normalized valuation, it holds also for a discrete valuation.

Definition 7. The field $\mathcal{K} := \mathcal{O}/\mathfrak{p}$ is called the **residue field** of \mathcal{O} .

We have an interesting property when a valuation is normalized.

Lemma 3. Let v be a normalized valuation on K , then for all $0 \neq x \in K$ we can write $x = ut^n$, where $t \in \mathfrak{p}$ with $v(t) = 1$, $u \in \mathcal{O}^*$ and $n \in \mathbb{Z}$. An element $x \in \mathfrak{p}$ such that $v(x) = 1$ is called **prime element**.

Proof: Since $v(K^*) = \mathbb{Z}$, there exists an element $t \in K$ with $v(t) = 1$. Therefore $t \in \mathfrak{p}$. Let $0 \neq x \in K$. We have that $v(x) = m$ for some $m \in \mathbb{Z}$. Hence $v(xt^{-m}) = 0$ and so $u := xt^{-m} \in \mathcal{O}^*$ and finally $x = ut^m$.

□

Recall that an ideal of a ring is principal, if it is generated by one element.

Theorem 4. Assume v is a normalized valuation on K , then \mathcal{O} is a principal ideal domain, i.e., every ideal of \mathcal{O} is principal. Moreover, all the ideals of \mathcal{O} different from $\{0\}$ are of the form

$$\mathfrak{p}^n = t^n \mathcal{O} = \{x \in K \mid v(x) \geq n\}, \quad n \geq 0,$$

where $t \in \mathfrak{p}$, $v(t) = 1$. Furthermore, we have

$$\mathfrak{p}^n / \mathfrak{p}^{n+1} \cong \mathcal{K}, \quad n \geq 1,$$

as K -vector spaces.

Proof: Let $\{0\} \neq A \subseteq \mathcal{O}$ be an ideal of the valuation ring and $0 \neq x \in A$ such that $n := v(x) \leq v(y)$ for all $y \in A$. By lemma 3, $x = ut^n$ for some prime element $t \in \mathfrak{p}$ and some $u \in \mathcal{O}^*$. This implies that $t^n \mathcal{O} \subseteq A$. Take now $y \in A$. According to lemma 3, $y = wt^m$, $w \in \mathcal{O}^*$. Since $y \in A$, $m = v(y) \geq n = v(x)$, so we can write $y = (wt^{m-n})t^n \in t^n \mathcal{O}$, hence $A \subseteq t^n \mathcal{O}$.

Consider now the K -homomorphism

$$\begin{aligned} \phi : \mathfrak{p}^n &\longrightarrow \mathcal{O}/\mathfrak{p} \\ at^n &\longmapsto a \pmod{\mathfrak{p}}, \end{aligned}$$

with $t \in \mathfrak{p}$ a prime element and $a \in \mathcal{O}$. Clearly, ϕ is surjective and its kernel is \mathfrak{p}^{n+1} . Hence, ϕ is a K -isomorphism.

□

This theorem gives us a useful lemma.

Lemma 4. *Let v, v' be two discrete valuation on K such that $\mathcal{O}_v = \mathcal{O}_{v'}$, where \mathcal{O}_v , resp. $\mathcal{O}_{v'}$, denotes the valuation ring generated by v , resp. by v' . Then v and v' are equivalent.*

Proof: Clearly, $\mathcal{O}_v = \mathcal{O}_{v'}$ implies $\mathcal{O}_v^* = \mathcal{O}_{v'}^*$ and $\mathfrak{p}_v = \mathfrak{p}_{v'}$. By theorem 4, there is an element $t \in \mathfrak{p}_v = \mathfrak{p}_{v'}$ such that $\mathfrak{p}_v = t\mathcal{O}_v = t\mathcal{O}_{v'} = \mathfrak{p}_{v'}$. Again by theorem 4, t must have the minimal valuation among the elements of $\mathfrak{p}_v = \mathfrak{p}_{v'}$. Set $v(t) =: s$ and $v'(t) =: s'$ and this gives $v(K^*) = s\mathbb{Z}$ and $v'(K^*) = s'\mathbb{Z}$. Clearly, for all $x \in \mathcal{O}_v^* = \mathcal{O}_{v'}^*$, $0 = v(x) = v'(x) = 0$. Consider $x \in K \setminus \mathcal{O}_v^* = K \setminus \mathcal{O}_{v'}^*$. Then, by lemma 3, $x = t^n u$ for some $n \in \mathbb{Z}$ and some $u \in \mathcal{O}_v^* = \mathcal{O}_{v'}^*$. So, $v(x) = ns$ and $v'(x) = s'n$ and this gives $v(x) = \frac{s}{s'}v'(x)$. Since x was arbitrary, the result holds for all $x \in K \setminus \mathcal{O}_v^* = K \setminus \mathcal{O}_{v'}^*$. Therefore v and v' are equivalent. □

1.5 Example: the field of rational numbers \mathbb{Q}

In this paragraph, we deal with the field of rational numbers \mathbb{Q} .

We know that on \mathbb{Q} we have the restriction of the Archimedean absolute value on \mathbb{R}

$$\begin{aligned} | \cdot | : \mathbb{Q} &\longrightarrow \mathbb{R} \\ x &\longmapsto \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases} . \end{aligned}$$

Since $| \cdot |$ is Archimedean, all the constructions that we made above are worthless.

We want a non-Archimedean absolute value on \mathbb{Q} . Consider a prime $p > 1$. For all $x \in \mathbb{Q}$ we can write

$$x = p^n \frac{a}{b}$$

with $n \in \mathbb{Z}$, $a \in \mathbb{Z}$, $0 \neq b \in \mathbb{N}$, $(a, b) = 1$, $p \nmid a$ and $p \nmid b$. Define the function $v_p : \mathbb{Q} \rightarrow \mathbb{R} \cup \{\infty\}$ as follows:

$$v_p(x) = v_p\left(p^n \frac{a}{b}\right) = n,$$

for all $0 \neq x \in \mathbb{Q}$, and $v_p(0) = \infty$. Clearly v_p is normalized valuation. According to theorem 3, we have the valuation ring

$$\mathcal{O}_p = \left\{ \frac{a}{b} \mid a \in \mathbb{Z}, b \in \mathbb{N} \setminus \{0\}, (a, b) = 1, p \nmid b \right\},$$

its group of units

$$\mathcal{O}_p^* = \left\{ \frac{a}{b} \mid a \in \mathbb{Z}, b \in \mathbb{N} \setminus \{0\}, (a, b) = 1, p \nmid a, p \nmid b \right\},$$

and its unique maximal ideal

$$\mathfrak{p}_p = \left\{ \frac{a}{b} \mid a \in \mathbb{Z}, b \in \mathbb{N} \setminus \{0\}, (a, b) = 1, p \mid a, p \nmid b \right\}.$$

Clearly $p \in \mathbb{Q}$ is a prime element and therefore all the non-zero ideals of \mathcal{O}_p are of the form

$$\mathfrak{p}_p^n = p^n \mathcal{O}_p, \quad n \geq 0.$$

Let now $x = p^n \frac{a}{b} \in \mathcal{O}_p$, $(a, b) = 1$, $a \in \mathbb{Z}$, $b \in \mathbb{N} \setminus \{0\}$, $p \nmid a$, $p \nmid b$ and $n \in \mathbb{N}$. If $n > 0$, $x \in \mathfrak{p}_p$ and therefore $x \equiv 0 \pmod{\mathfrak{p}_p}$. Suppose $n = 0$, then $x \in \mathcal{O}_p^*$. We now that for every $m \in \mathbb{Z}$, we can write

$$m = \sum_{i=0}^r m_i p^i$$

with $r, m_i \in \mathbb{N}$, $0 \leq m_i < p$, $0 \leq i \leq r$. Hence, we have

$$a = \sum_{i=0}^s a_i p^i \quad \text{and} \quad b = \sum_{j=0}^t b_j p^j$$

with $s, t, a_i, b_j \in \mathbb{N}$, $0 \leq a_i, b_j < p$, $0 \leq i \leq s$, $0 \leq j \leq t$. This gives

$$\begin{aligned} x &= \frac{a}{b} \\ &= \frac{a_0 + a_1 p + \cdots + a_s p^s}{b} \\ &= \frac{a_0}{b} + \frac{a_1 p + \cdots + a_s p^s}{b} \end{aligned}$$

with

$$\frac{a_1 p + \cdots + a_s p^s}{b} = p \frac{a_1 + \cdots + a_s p^{s-1}}{b} \in \mathfrak{p}_p.$$

Moreover, we have that

$$\frac{a_0}{b} = \frac{a_0}{b_0 + b_1 p + \cdots + b_t p^t} = \frac{a_0}{b_0} - \frac{a_0 b_1 p + a_0 b_2 p^2 + \cdots + a_0 b_t p^t}{b_0^2 + b_0 b_1 p + \cdots + b_0 b_t p^t}$$

and so

$$\frac{a_0 b_1 p + a_0 b_2 p^2 + \cdots + a_0 b_t p^t}{b_0^2 + b_0 b_1 p + \cdots + b_0 b_t p^t} \in \mathfrak{p}_p.$$

This means that

$$x \equiv \frac{a_0}{b_0} \pmod{\mathfrak{p}_p}$$

and finally we get that

$$\mathbb{F}_p = \mathbb{Z}/p\mathbb{Z} \cong \mathcal{K}_p = \mathcal{O}_p/\mathfrak{p}_p.$$

By theorem 2, we know that each valuation v_p gives a corresponding family of equivalent non-Archimedean absolute values. We will denote by $|\cdot|_p$ the absolute value

$$|x|_p := p^{-v_p(x)},$$

for all $0 \neq x \in \mathbb{Q}$.

We conclude this example with an important result.

Theorem 5. On \mathbb{Q} , each non-trivial absolute value is equivalent to an absolute value $|\cdot|_p$ or $|\cdot|$.

Proof: Let $\|\cdot\|$ be a non-Archimedean absolute value on \mathbb{Q} . We have

$$\|n\| = \|1 + \cdots + 1\| \leq \max\{\|1\|, \dots, \|1\|\} = 1.$$

There exists a prime $p > 1$ such that $\|p\| < 1$. If not, $\|\cdot\|$ is the trivial valuation. Consider the set

$$A := \{a \in \mathbb{Z} \mid \|a\| < 1\}.$$

Clearly, A is an ideal of \mathbb{Z} and $p\mathbb{Z} \subseteq A \neq \mathbb{Z}$. Since $p\mathbb{Z}$ is a maximal ideal, we have that $A = p\mathbb{Z}$. Take $a \in \mathbb{Z}$, $a = p^m b$ with $p \nmid b$. This implies that $b \notin A$ and therefore $\|b\| = 1$. This gives

$$\|a\| = \|p^m b\| = \|p\|^m = |a|_p^s,$$

with $s := -\frac{\log\|p\|}{\log p}$. Therefore $\|\cdot\|$ is equivalent to $|\cdot|_p$.

Assume now that $\|\cdot\|$ is Archimedean. For all integer $m, n > 1$, we have

$$\|m\|^{\frac{1}{\log m}} = \|n\|^{\frac{1}{\log n}}.$$

Indeed, we can write

$$m = a_0 + a_1 n + \cdots + a_r n^r$$

with $r, a_i \in \mathbb{N}$, $0 \leq a_i < n$, $0 \leq i \leq r$. Clearly $n^r \leq m$, and so $r \leq \frac{\log m}{\log n}$. Moreover,

$$\|a_i\| \leq \|1 + \cdots + 1\| \leq a_i \|1\| \leq n$$

gives that

$$\|m\| \leq \sum_{i=0}^r \|a_i\| \|n\|^i \leq \sum_{i=0}^r \|a_i\| \|n\|^r \leq \left(1 + \frac{\log m}{\log n}\right) n \|n\|^{\frac{\log m}{\log n}}.$$

Replacing m by m^k , $k \in \mathbb{N}$, and taking the k -th root, we get that

$$\|m\| = \sqrt[k]{\|m^k\|} \leq \sqrt[k]{\left(1 + \frac{k \log m}{\log n}\right) n \|n\|^{\frac{k \log m}{\log n}}} = \|n\|^{\frac{\log m}{\log n}} \sqrt[k]{\left(1 + \frac{k \log m}{\log n}\right) n},$$

and when k goes to ∞ we have

$$\|m\| \leq \|n\|^{\frac{\log m}{\log n}} \text{ and so } \|m\|^{\frac{1}{\log m}} \leq \|n\|^{\frac{1}{\log n}}.$$

Exchanging the roles of m and n we get the inverse inequality.

Since $\|n\|^{\frac{1}{\log n}} > 0$, there exists $s \in \mathbb{R}$ such that $e^s = \|n\|^{\frac{1}{\log n}}$ and so $\|n\| = e^{s \log n}$. Hence, for all $x \in \mathbb{Q}$, $x > 0$, we have

$$\|x\| = e^{s \log x} = x^s = |x|^s$$

Since $\|x\| = \|-x\|$, we have that $\|\cdot\|$ is equivalent to $|\cdot|$.

□

1.6 Example: the rational function field $\mathbb{F}_q(T)$

Let $q = p^n$, $p > 1$ a prime and $n \in \mathbb{N} \setminus \{0\}$. The field

$$\mathbb{F}_q(T) = \left\{ \frac{f(T)}{g(T)} \mid f(T) \in \mathbb{F}_q[T], g(T) \in \mathbb{F}_q[T] \setminus \{0\} \right\}$$

is called **rational function field**.

A polynomial is called **monic** if the leading coefficient is equal to 1. Let $p(T) \in \mathbb{F}_q[T]$ be a monic, irreducible polynomial. For all $f(T) \in \mathbb{F}_q(T)$ we can write

$$f(T) = p(T)^n \frac{g(T)}{h(T)},$$

with $n \in \mathbb{Z}$, $g(T) \in \mathbb{F}_q[T]$ such that $p(T) \nmid g(T)$ and $h(T) \in \mathbb{F}_q[T] \setminus \{0\}$ such that $p(T) \nmid h(T)$.

Define the function $v_{p(T)} : \mathbb{F}_q(T) \rightarrow \mathbb{Z} \cup \{\infty\}$ as follows:

$$v_{p(T)}(f(T)) = v_{p(T)} \left(p(T)^n \frac{g(T)}{h(T)} \right) := n,$$

for all $0 \neq f(T) \in \mathbb{F}_q(T)$ and $v_{p(T)}(0) = \infty$. Obviously, $v_{p(T)}$ is a normalized valuation and we have, according to theorem 3, the valuation ring

$$\mathcal{O}_{p(T)} = \left\{ \frac{f(T)}{g(T)} \mid f(T) \in \mathbb{F}_q[T], g(T) \in \mathbb{F}_q[T] \setminus \{0\}, p(T) \nmid g(T) \right\},$$

its group of units

$$\mathcal{O}_{p(T)}^* = \left\{ \frac{f(T)}{g(T)} \mid f(T) \in \mathbb{F}_q[T], g(T) \in \mathbb{F}_q[T] \setminus \{0\}, p(T) \nmid f(T), p(T) \nmid g(T) \right\},$$

and its unique maximal ideal

$$\mathfrak{p}_{p(T)} = \left\{ \frac{f(T)}{g(T)} \mid f(T) \in \mathbb{F}_q[T], g(T) \in \mathbb{F}_q[T] \setminus \{0\}, p(T) \mid f(T), p(T) \nmid g(T) \right\}.$$

Moreover, the residue field of $\mathcal{O}_{p(T)}$, $\mathcal{K}_{p(T)} = \mathcal{O}_{p(T)}/\mathfrak{p}_{p(T)}$ is isomorphic to $\mathbb{F}_q[T]/(p(T))$. Indeed, consider the ring homomorphism

$$\begin{aligned} \varphi : \mathbb{F}_q[T] &\longrightarrow \mathcal{O}_{p(T)}/\mathfrak{p}_{p(T)} \\ f(T) &\longmapsto f(T) \pmod{\mathfrak{p}_{p(T)}}. \end{aligned}$$

Clearly, the kernel of φ is the ideal $(p(T))$ generated by $p(T)$ in $\mathbb{F}_q[T]$. Take now $h(T) \in \mathcal{O}_{p(T)}$. We can write $h(T) = \frac{r(T)}{s(T)}$ with $r(T), s(T) \in \mathbb{F}_q[T]$, such that $s(T) \neq 0$ and $p(T) \nmid s(T)$. Thus, there exist $a(T), b(T) \in \mathbb{F}_q[T]$ with $a(T)p(T) + b(T)s(T) = 1$ and therefore

$$h(T) = 1 \cdot h(T) = \frac{a(T)r(T)}{s(T)}p(T) + b(T)r(T),$$

and so

$$h(T) \equiv b(T)r(T) \pmod{\mathfrak{p}_{p(T)}}.$$

Since $b(T), r(T) \in \mathbb{F}_q[T]$, φ is surjective and we have an isomorphism

$$\mathbb{F}_q[T]/(p(T)) \cong \mathcal{K}_{p(T)} = \mathcal{O}_{p(T)}/\mathfrak{p}_{p(T)}.$$

Let $f(T) \in \mathbb{F}_q(T)$. We can write

$$f(T) = \frac{g(T)}{h(T)},$$

with $g(T) \in \mathbb{F}_q[T]$, $h(T) \in \mathbb{F}_q[T] \setminus \{0\}$. Consider now the function

$$v_\infty : \mathbb{F}_q(T) \rightarrow \mathbb{Z} \cup \{\infty\}$$

defined as follows:

$$v_\infty(f(T)) = v_\infty\left(\frac{g(T)}{h(T)}\right) := \deg h(T) - \deg g(T),$$

for all $0 \neq f(T) \in \mathbb{F}_q(T)$ and $v_\infty(0) = \infty$. Obviously, v_∞ is a normalized valuation and, by theorem 3, we have the valuation ring

$$\mathcal{O}_\infty = \left\{ \frac{f(T)}{g(T)} \mid f(T) \in \mathbb{F}_q[T], g(T) \in \mathbb{F}_q[T] \setminus \{0\}, \deg f(T) \leq \deg g(T) \right\},$$

its group of units

$$\mathcal{O}_\infty^* = \left\{ \frac{f(T)}{g(T)} \mid f(T) \in \mathbb{F}_q[T], g(T) \in \mathbb{F}_q[T] \setminus \{0\}, \deg f(T) = \deg g(T) \right\},$$

and its unique maximal ideal

$$\mathfrak{p}_\infty = \left\{ \frac{f(T)}{g(T)} \mid f(T) \in \mathbb{F}_q[T], g(T) \in \mathbb{F}_q[T] \setminus \{0\}, \deg f(T) < \deg g(T) \right\}.$$

We have that $T^{-1} \in \mathbb{F}_q(T)$ is a prime element, since $v_\infty(T^{-1}) = \deg T - \deg 1 = 1$. Therefore, all non-zero ideal are of the form

$$\mathfrak{p}_\infty^n = (T)^{-n} \mathcal{O}_\infty, \quad n \geq 0.$$

Take $f(T) \in \mathcal{O}_\infty$, $f(T) = \frac{g(T)}{h(T)}$, with $g(T) \in \mathbb{F}_q[T]$, $h(T) \in \mathbb{F}_q[T] \setminus \{0\}$ and $n := \deg g(T) \leq \deg h(T) =: m$. Then we can write

$$g(T) = a_n T^n + a_{n-1} T^{n-1} + \cdots + a_0 \quad \text{and} \quad h(T) = b_m T^m + b_{m-1} T^{m-1} + \cdots + b_0,$$

with $a_i \in \mathbb{F}_q$, $0 \leq i \leq n$, $a_n \neq 0$, and $b_j \in \mathbb{F}_q$, $0 \leq j \leq m$, $b_m \neq 0$. If $n < m$, we have $f(T) \equiv 0 \pmod{\mathfrak{p}_\infty}$. If $n = m$, we have

$$\begin{aligned} f(T) &= \frac{g(T)}{h(T)} \\ &= \frac{a_n T^n + a_{n-1} T^{n-1} + \cdots + a_0}{b_n T^n + b_{n-1} T^{n-1} + \cdots + b_0} \\ &= \frac{a_n T^n}{b_n T^n + b_{n-1} T^{n-1} + \cdots + b_0} + \frac{a_{n-1} T^{n-1} + \cdots + a_0}{b_n T^n + b_{n-1} T^{n-1} + \cdots + b_0}, \end{aligned}$$

with

$$\frac{a_{n-1}T^{n-1} + \cdots + a_0}{b_nT^n + b_{n-1}T^{n-1} + \cdots + b_0} \in \mathfrak{p}_\infty.$$

Moreover

$$\frac{a_nT^n}{b_nT^n + b_{n-1}T^{n-1} + \cdots + b_0} = \frac{a_n}{b_n} - \frac{a_nb_{n-1}T^{n-1} + a_nb_{n-2}T^{n-2} + \cdots + a_nb_0}{b_n^2T^n + b_nb_{n-1}T^{n-1} + \cdots + b_nb_0},$$

with

$$\frac{a_nb_{n-1}T^{n-1} + a_nb_{n-2}T^{n-2} + \cdots + a_nb_0}{b_n^2T^n + b_nb_{n-1}T^{n-1} + \cdots + b_nb_0} \in \mathfrak{p}_\infty.$$

Therefore, we get that

$$f(T) \equiv \frac{a_n}{b_n} \pmod{\mathfrak{p}_\infty}.$$

This means that

$$\mathcal{K}_\infty \cong \mathbb{F}_q.$$

Note that in the case of the above valuations, we can embed the field \mathbb{F}_q in the residue field and this leads to an important

Definition 8. *Let v be a valuation on $\mathbb{F}_q(T)$ and \mathfrak{p}_v the maximal ideal of the valuation ring with respect to v . Then*

$$\deg \mathfrak{p}_v := [\mathcal{K}_v : \mathbb{F}_q]$$

*is called the **degree** of \mathfrak{p}_v .*

Note that if a valuation is defined as above by an irreducible, monic polynomial $p(T) \in \mathbb{F}_q[T]$, the degree of $\mathfrak{p}_{p(T)}$ is $\deg \mathfrak{p}_{p(T)} = \deg p(T)$. Indeed $\mathcal{K}_{p(T)} \cong \mathbb{F}_q[T]/(p(T))$ and therefore $\deg \mathfrak{p}_{p(T)} = [\mathcal{K}_{p(T)} : \mathbb{F}_q] = \deg p(T)$. Moreover $\deg \mathfrak{p}_\infty = 1$, since $\mathcal{K}_\infty \cong \mathbb{F}_q$.

We have an interesting lemma about the degree.

Lemma 5. *Let v be a valuation on $\mathbb{F}_q(T)$ and $0 \neq f(T) \in \mathfrak{p}_v$. Then we have*

$$\deg \mathfrak{p}_v \leq [\mathbb{F}_q(T) : \mathbb{F}_q(f(T))] < \infty,$$

where $\mathbb{F}_q(f(T))$ denotes the field generated by $f(T)$ over \mathbb{F}_q .

Proof: Note that $\mathbb{F}_q \subset \mathbb{F}_q(f(T))$. Write $f(T) = \frac{g(T)}{h(T)}$, with $g(T), h(T) \in \mathbb{F}_q[T]$. Then, $p(X) := f(T)h(X) - g(X)$ is a polynomial in X with coefficients in $\mathbb{F}_q(f(T))$. Clearly T is a root of $p(X)$, hence $[\mathbb{F}_q(T) : \mathbb{F}_q(f(T))] < \infty$. For the remaining inequality, it suffices to show that any $f_1(T), \dots, f_n(T) \in \mathcal{O}_v$, whose residue class in $\bar{f}_1(T), \dots, \bar{f}_n(T) \in \mathcal{K}_v$ are linearly independent over \mathbb{F}_q , are linearly independent over $\mathbb{F}_q(f(T))$. Assume that

$$\sum_{i=0}^n \varphi_i f_i(T) = 0$$

for some $\varphi_i \in \mathbb{F}_q(f(T))$, $0 \leq i \leq n$, not all zero. In fact, φ_i are linear combinations of powers of $f(T)$ with coefficients on \mathbb{F}_q . W.l.o.g. we can assume that φ_i are polynomials in $f(T)$ and not all of the φ_i are divisible by $f(T)$, so we can write $\varphi_i = a_i + f(T)g_i$, with $a_i \in \mathbb{F}_q$ not all zero, $g_i \in \mathbb{F}_q[f(T)]$, $0 \leq i \leq n$. Since $f(T) \in \mathfrak{p}_v$ and $g_i \in \mathbb{F}_q[f(T)] \subseteq \mathcal{O}_v$, we have that $\varphi_i \equiv a_i \pmod{\mathfrak{p}_v}$, for all $0 \leq i \leq n$. Hence

$$0 \equiv \sum_{i=0}^n \varphi_i f_i(T) \equiv \sum_{i=0}^n a_i \bar{f}_i(T) \pmod{\mathfrak{p}_v}.$$

This contradicts the linear independence of $\bar{f}_1(T), \dots, \bar{f}_n(T) \in \mathcal{K}_v$ over \mathbb{F}_q .

□

We can state now a very important theorem concerning the rational function field $\mathbb{F}_q(T)$.

Theorem 6. *All the non-trivial valuations on the rational function field $\mathbb{F}_q(T)$ are equivalent to a valuation $v_{p(T)}$, for $p(T) \in \mathbb{F}_q[T]$ a monic, irreducible polynomial, or to v_∞ .*

Proof: By lemma 4, it suffices to show that if v is a non-trivial valuation different from v_∞ , there is an irreducible, monic polynomial $p(T) \in \mathbb{F}_q[T]$ such that

$$\mathcal{O}_{p(T)} = \mathcal{O}_v.$$

Assume first that $T \in \mathcal{O}_v$, then, obviously, $\mathbb{F}_q[T] \subseteq \mathcal{O}_v$. Set $I := \mathbb{F}_q[T] \cap \mathfrak{p}_v$; this is a prime ideal of $\mathbb{F}_q[T]$. Indeed \mathfrak{p}_v is a maximal ideal, therefore is prime, and $\mathbb{F}_q[T]$ is a ring. The map $\mathbb{F}_q[T] \rightarrow \mathcal{K}_v$ induces an embedding $\mathbb{F}_q[T]/I \hookrightarrow \mathcal{K}_v$, and therefore, by lemma 5, $I \neq \{0\}$. Indeed, if $I = \{0\}$, we have that $\mathbb{F}_q[T]/I = \mathbb{F}_q[T]$ and so $\mathbb{F}_q[T] \hookrightarrow \mathcal{K}_v$. But $\infty = [\mathbb{F}_q[T] : \mathbb{F}_q] \leq [\mathcal{K}_v : \mathbb{F}_q] < \infty$. Since $I \neq \{0\}$ and I is prime, there is a unique irreducible, monic polynomial $p(T) \in \mathbb{F}_q[T]$ such that $I = p(T)\mathbb{F}_q[T]$ (recall that the ring $\mathbb{F}_q[T]$ is principal). Any $g(T) \in \mathbb{F}_q[T]$ with $p(T) \nmid g(T)$ is not in I , so $g(T) \notin \mathfrak{p}_v \subset \mathcal{O}_v$, therefore $g(T)^{-1} \in \mathcal{O}_v$. Hence, we have

$$\mathcal{O}_{p(T)} = \left\{ \frac{f(T)}{g(T)} \mid f(T) \in \mathbb{F}_q[T], g(T) \in \mathbb{F}_q[T] \setminus \{0\}, p(T) \nmid g(T) \right\} \subseteq \mathcal{O}_v.$$

By theorem 3, all valuation rings are maximal proper subrings of $\mathbb{F}_q(T)$, therefore $\mathcal{O}_{p(T)} = \mathcal{O}_v$. Therefore v and $v_{p(T)}$ are equivalent.

Assume now that $T \notin \mathcal{O}_v$. We have that $\mathbb{F}_q[T^{-1}] \subseteq \mathcal{O}_v$, $T^{-1} \in \mathfrak{p}_v \cap \mathbb{F}_q[T^{-1}]$ and clearly $\mathfrak{p}_v \cap \mathbb{F}_q[T^{-1}] = T^{-1}\mathbb{F}_q[T^{-1}]$, since $\mathfrak{p}_v \cap \mathbb{F}_q[T^{-1}]$ is a prime ideal. As before, if $g(T^{-1}) \in \mathbb{F}_q[T^{-1}]$ with $T^{-1} \nmid g(T^{-1})$, $g(T^{-1}) \notin \mathfrak{p}_v$ and so $g(T^{-1})^{-1} \in \mathcal{O}_v$. This

gives

$$\begin{aligned}
\mathcal{O}_v &\supseteq \left\{ \frac{f(T^{-1})}{g(T^{-1})} \mid f(T^{-1}) \in \mathbb{F}_q[T^{-1}], g(T^{-1}) \in \mathbb{F}_q[T^{-1}] \setminus \{0\}, T^{-1} \nmid g(T^{-1}) \right\} \\
&= \left\{ \frac{a_0 + a_1 T^{-1} + \dots + a_n T^{-n}}{b_0 + b_1 T^{-1} + \dots + b_m T^{-m}} \mid b_0 \neq 0 \right\} \\
&= \left\{ \frac{a_0 T^{m+n} + a_1 T^{m+n-1} + \dots + a_n T^m}{b_0 T^{m+n} + b_1 T^{m+n-1} + \dots + b_m T^n} \mid b_0 \neq 0 \right\} \\
&= \left\{ \frac{u(T)}{v(T)} \mid u(T) \in \mathbb{F}_q[T], v(T) \in \mathbb{F}_q[T] \setminus \{0\}, \deg u(T) \leq \deg v(T) \right\} \\
&= \mathcal{O}_\infty.
\end{aligned}$$

According to theorem 3, this implies that $\mathcal{O}_v = \mathcal{O}_\infty$ and so v is equivalent to v_∞ .

□

This theorem and the theorem 5 for \mathbb{Q} are very similar. The valuations v_p , given by a prime p , on \mathbb{Q} correspond to the valuations $v_{p(T)}$, given by a monic, irreducible polynomial $p(T)$, on $\mathbb{F}_q(T)$ and we have a difference between the usual absolute value $|\cdot|$ on \mathbb{Q} and the valuation v_∞ on $\mathbb{F}_q(T)$. The problem is that $|\cdot|$ is Archimedean, hence we haven't any corresponding valuation.

We conclude this example with a corollary.

Corollary. *There is a bijection between the equivalence classes of non-trivial valuations v on $\mathbb{F}_q(T)$ with $\deg \mathfrak{p}_v = 1$ and $\mathbb{F}_q \cup \{\infty\}$.*

Proof: By theorem 6, the valuations $v_{p(T)}$, $p(T)$ an irreducible, monic polynomial in $\mathbb{F}_q[T]$, and v_∞ represent all the equivalence classes of non-trivial valuations on $\mathbb{F}_q(T)$. Moreover, by the note after the definition 8, the valuations with degree 1 are exactly v_∞ and $v_{T-\alpha}$, for all $\alpha \in \mathbb{F}_q$. Therefore, we have a bijection between $\mathbb{F}_q \cup \{\infty\}$ and the equivalence classes of non-trivial valuations on $\mathbb{F}_q(T)$.

□

Remark: In fact all the results that holds for $\mathbb{F}_q(T)$, holds also for all fields

$$K(T) := \left\{ \frac{f(T)}{g(T)} \mid f(T) \in K[T], g(T) \in K[T] \setminus \{0\} \right\},$$

where $K[T]$ denotes the ring of polynomials over a field K . $K(T)$ is called **rational function field**.

2 Completion

2.1 Definitions and results

We begin this section with some basic definitions.

Definition 9. Let K be a field and $|\cdot|$ an absolute value on K . A sequence $\{a_n\}_{n \in \mathbb{N}}$ in K is called a **Cauchy sequence** if, for all $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$n, m \geq N \implies |a_n - a_m| < \varepsilon.$$

Another important

Definition 10. A field K with an absolute value $|\cdot|$ is called **complete** if any Cauchy sequence $\{a_n\}_{n \in \mathbb{N}}$ in K converges to an element $a \in K$, i.e.

$$\lim_{n \rightarrow \infty} |a_n - a| = 0.$$

A useful lemma for non-Archimedean absolute values.

Lemma 6. Let K be a complete field and $|\cdot|$ a non-Archimedean absolute value on K . Then, for $\{a_n\}_{n \in \mathbb{N}} \subset K$, we have:

1. the sequence $\{a_n\}_{n \in \mathbb{N}}$ is a Cauchy sequence if, and only if, $\lim_{n \rightarrow \infty} (a_{n+1} - a_n) = 0$,
2. the series $\sum_{n=0}^{\infty} a_n$ converges if, and only if, $\lim_{n \rightarrow \infty} a_n = 0$,
3. Suppose that $\lim_{n \rightarrow \infty} a_n = a \neq 0$, then there exists a positive integer N such that for all $m \geq N$, $|a_m| = |a_N| = |a|$.

Proof:

1. Assume that $\{a_n\}_{n \in \mathbb{N}}$ is a Cauchy sequence, then, for all $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that for all $m, n \geq N$, $|a_m - a_n| < \varepsilon$, therefore we have also, for all $n \geq N$, $|a_{n+1} - a_n| < \varepsilon$. Conversely, assume that for all $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that for all $n \geq N$, $|a_{n+1} - a_n| < \varepsilon$. Then, for all $r, s \geq N$,

$$\begin{aligned} |a_r - a_s| &= \left| \sum_{i=n}^{m-1} (a_{i+1} - a_i) \right| \\ &\leq \max_{n \leq i < m} \{|a_{i+1} - a_i|\} < \varepsilon. \end{aligned}$$

Therefore, $\{a_n\}_{n \in \mathbb{N}}$ is a Cauchy sequence.

2. We have that $|a_n| = \left| \sum_{i=0}^n a_i - \sum_{j=0}^{n-1} a_j \right|$. Using 1., the equivalence is obvious.

3. Since $\lim_{n \rightarrow \infty} a_n = a \neq 0$, there exists a positive integer n_a such that, for all $n \geq n_a$, $|a_n - a| < |a|$. Then, using the note we made after definition 3, we have that $|a_n| = |a_n - a + a| = \max\{|a_n - a|, |a|\} = |a|$.

□

Note that for a non-Archimedean absolute value we have the equivalence

$$\exists a > 0 \text{ such that } |x| < a \Leftrightarrow \exists b \in \mathbb{R} \text{ such that } v(x) > b,$$

with v a corresponding valuation. Indeed, for $s > 0$, if $|x| < a$, $v(x) = -s \log |x| > -s \log a$. Conversely, for $q > 1$, if $v(x) > b$, $|x| = q^{-v(x)} < q^{-b}$.

Theorem 7. *Let K be a field and $|\cdot|$ be an absolute value on K . Then, there exists a unique, up to K -isomorphism, complete field \widehat{K} with an absolute value $|\cdot|_{\widehat{K}}$ such that K is embedded in \widehat{K} as a dense subfield and the absolute value on K is a restriction of the absolute value on \widehat{K} , i.e., $|x|_{\widehat{K}} = |x|$ if $x \in K$.*

Sketch of the proof: We will not prove this theorem in detail, not because it is too difficult, but because we would need to prove a lot of uninteresting little claims, that can be easily proved by the reader.

Let R be the set of all the Cauchy sequences in K with respect to $|\cdot|$. Define the addition and the multiplication as follows, for all $\{a_n\}_{n \in \mathbb{N}}, \{b_n\}_{n \in \mathbb{N}} \in R$:

$$\{a_n\}_{n \in \mathbb{N}} + \{b_n\}_{n \in \mathbb{N}} := \{a_n + b_n\}_{n \in \mathbb{N}} \text{ and } \{a_n\}_{n \in \mathbb{N}} \cdot \{b_n\}_{n \in \mathbb{N}} := \{a_n b_n\}_{n \in \mathbb{N}}.$$

Indeed R is a ring. Let $\mathfrak{m} \subset R$ be the set of all the Cauchy sequences that converge to 0. It not difficult to prove that \mathfrak{m} is a maximal ideal of R . Now set

$$\widehat{K} := R/\mathfrak{m}.$$

Clearly, \widehat{K} is a field. We have an injection $K \hookrightarrow \widehat{K}$ by sending $a \in K$ to the equivalence class of the Cauchy sequence (a, a, a, \dots) . Hence, we can write $K \subset \widehat{K}$. Take $a \in \widehat{K}$ and let $\{a_n\}_{n \in \mathbb{N}} \in R$ be a representative of a . Then, we have that the sequence $\{|a_n|\}_{n \in \mathbb{N}}$ converges in \mathbb{R} , because it is a Cauchy sequence, since $||a_n| - |a_m|| \leq |a_n - a_m|$, by 4. of lemma 1. Set

$$|a|_{\widehat{K}} := \lim_{n \rightarrow \infty} |a_n|,$$

then $|\cdot|_{\widehat{K}}$ is an absolute value on \widehat{K} and, if $a \in K$, we have $|a|_{\widehat{K}} = |a|$. Furthermore,

$$\lim_{n \rightarrow \infty} a_n = a$$

in \widehat{K} , therefore K is dense in \widehat{K} and \widehat{K} is complete with respect to $|\cdot|_{\widehat{K}}$.

Let \widehat{K}' be another complete field, with respect to an absolute value $|\cdot|_{\widehat{K}'}$, such that K is dense in \widehat{K}' and, for all $x \in K$, $|x|_{\widehat{K}'} = |x|$. Take $a \in \widehat{K}$ and let $\{a_n\}_{n \in \mathbb{N}} \subset K$ be a representative of a . Then, in \widehat{K}' , this Cauchy sequence converges to an element $a' \in \widehat{K}'$, because K is dense in \widehat{K}' . Define the function $\sigma : \widehat{K} \rightarrow \widehat{K}'$ by $\sigma(a) := a'$. It is easy to verify that σ is a K -isomorphism. Furthermore, $|a|_{\widehat{K}} = |\sigma(a)|_{\widehat{K}'}$ because

$$|a|_{\widehat{K}} = \lim_{n \rightarrow \infty} |a_n| = \lim_{n \rightarrow \infty} |a_n|_{\widehat{K}'} = |a|_{\widehat{K}'}.$$

Definition 11. The field \widehat{K} is called the **completion** of K .

Now we look at an Archimedean absolute value on a field K . The following theorem is due to Alexander Ostrowski (1893-1986). We will not prove this theorem.

Theorem 8. Let K be a complete field with respect to an Archimedean absolute value $|\cdot|_K$, then there is an isomorphism σ from K to \mathbb{R} or \mathbb{C} such that $|x|_K = |\sigma(x)|^s$ for all $x \in K$ and a fixed $0 \leq s \leq 1$, where $|\cdot|$ denotes the usual absolute value of \mathbb{R} or \mathbb{C} .

Ostrowski's theorem tells us that all complete fields with respect to an Archimedean absolute value are isomorphic to \mathbb{R} or \mathbb{C} . Therefore, the completion of a field with an Archimedean absolute value is isomorphic to \mathbb{R} or \mathbb{C} .

Let us look now at the completion of a field with a non-Archimedean absolute value. As seen before, there is a valuation v corresponding to the non-Archimedean absolute value on K . In this case, we denote \widehat{v} the valuation of the completion \widehat{K} of K . Clearly, if v is discrete, resp. normalized, \widehat{v} is also discrete, resp. normalized.

Theorem 9. Let K be a field, \widehat{K} its completion with respect to the valuation v on K . Denote \widehat{v} the corresponding valuation on \widehat{K} , \mathcal{O} , resp. $\widehat{\mathcal{O}}$ the valuation ring of K , resp. \widehat{K} , \mathfrak{p} , resp. $\widehat{\mathfrak{p}}$, the maximal ideal of \mathcal{O} , resp. $\widehat{\mathcal{O}}$ and \mathcal{K} , resp. $\widehat{\mathcal{K}}$, the residue field of \mathcal{O} , resp. $\widehat{\mathcal{O}}$. Then

$$\mathcal{K} \cong \widehat{\mathcal{K}}$$

and, if v is discrete,

$$\mathcal{O}/\mathfrak{p}^n \cong \widehat{\mathcal{O}}/\widehat{\mathfrak{p}}^n, \quad n \geq 1.$$

Proof: By theorem 7, we have $K \subset \widehat{K}$, $\mathcal{O} \subset \widehat{\mathcal{O}}$ and $\mathfrak{p} \subset \widehat{\mathfrak{p}}$. The inclusion $\mathcal{O} \subset \widehat{\mathcal{O}}$ gives a homomorphism

$$\varphi : \mathcal{O} \rightarrow \widehat{\mathcal{O}}/\widehat{\mathfrak{p}},$$

whose kernel is obviously \mathfrak{p} . Let now $x \in \widehat{\mathcal{O}}$. By theorem 7, K is dense in \widehat{K} , therefore there is a sequence $\{x_n\}_{n \in \mathbb{N}} \subset K$, which converges to $x \in \widehat{K}$. Since $\widehat{v}(x) \geq 0$, by lemma 6, there exists a positive integer N such that $\widehat{v}(x_n) = \widehat{v}(x)$, for all $n \geq N$. Hence, we can assume that $\{x_n\}_{n \in \mathbb{N}} \subset \mathcal{O}$. By definition, for all $\xi \in \mathbb{R}$, there is $N \in \mathbb{N}$ such that, for all $n \geq N$, $\widehat{v}(x - x_n) > \xi$. Take $\xi > 0$, then $x - x_n \in \widehat{\mathfrak{p}}$, and we get $x \equiv x_n \pmod{\widehat{\mathfrak{p}}}$. This means that φ is surjective and therefore we have an isomorphism

$$\mathcal{O}/\mathfrak{p} \cong \widehat{\mathcal{O}}/\widehat{\mathfrak{p}}.$$

Moreover, if v is discrete, \widehat{v} is discrete and all the ideal of \mathcal{O} , resp. $\widehat{\mathcal{O}}$, are of the form \mathfrak{p}^n , resp. $\widehat{\mathfrak{p}}^n$, $n \geq 1$. So we have a homomorphism $\lambda : \mathcal{O} \rightarrow \widehat{\mathcal{O}}/\widehat{\mathfrak{p}}^n$, whose kernel is \mathfrak{p}^n . By the same argument as above, for all $x \in \widehat{\mathcal{O}}$, there is an element $y_n \in \mathcal{O}$ such that $\widehat{v}(x - y_n) \geq n$, for all $n \geq 1$. Therefore $x \equiv y_n \pmod{\widehat{\mathfrak{p}}^n}$. Hence, λ is surjective and we have an isomorphism $\mathcal{O}/\mathfrak{p}^n \cong \widehat{\mathcal{O}}/\widehat{\mathfrak{p}}^n$.

□

Theorem 10. *Take the same assumption as in the preceding theorem and assume that v is normalized. Let $R \subseteq \mathcal{O}$ be a set of representatives of \mathcal{K} such that $0 \in R$ and let $t \in \mathfrak{p}$ be a prime element. Then we can represent all $x \in \widehat{K}^*$ as a converging series*

$$x = t^m(a_0 + a_1t + a_2t^2 + \dots)$$

with $a_i \in R$, $i \in \mathbb{N}$, $a_0 \neq 0$ and $m \in \mathbb{Z}$.

Proof: Since $\mathfrak{p} \subset \widehat{\mathfrak{p}}$, $t \in \widehat{\mathfrak{p}}$ and $1 = v(t) = \widehat{v}(t)$, according to theorem 7. From now on, in this proof, we will use an absolute value corresponding to the valuation \widehat{v} . By lemma 3, we have that $x = ut^m$, $u \in \widehat{\mathcal{O}}^*$. Since $\mathcal{O}/\mathfrak{p} \cong \widehat{\mathcal{O}}/\widehat{\mathfrak{p}}$, $u \pmod{\widehat{\mathfrak{p}}}$ has a representative $0 \neq a_0 \in R$ and therefore we can write $u = a_0 + tb_1$ with $b_1 \in \widehat{\mathcal{O}}$. By the same argument, we find also $a_1, a_2, \dots, a_{n-1} \in R$ such that

$$u = a_0 + a_1t + \dots + a_{n-1}t^{n-1} + t^n b_n$$

with $b_n \in \widehat{\mathcal{O}}$. As before, there is an $a_n \in R$ such that $b_n = a_n + tb_{n+1}$, $b_{n+1} \in \widehat{\mathcal{O}}$. Hence,

$$u = a_0 + a_1t + \dots + a_{n-1}t^{n-1} + a_nt^n + t^{n+1}b_{n+1}.$$

We can do this for all $n \in \mathbb{N}$, therefore we have a series

$$\sum_{r=0}^{\infty} a_r t^r.$$

It remains to show that this series converges to u . For all $n \in \mathbb{N}$, we have

$$\begin{aligned} \widehat{v}\left(u - \sum_{i=0}^n a_i t^i\right) &= \widehat{v}(t^{n+1}b_{n+1}) \\ &= \widehat{v}(t^{n+1}) + \widehat{v}(b_{n+1}) \\ &= n + 1 + \widehat{v}(b_{n+1}) \\ &\geq n + 1, \end{aligned}$$

since $b_{n+1} \in \widehat{\mathcal{O}}$. This gives

$$\lim_{n \rightarrow \infty} \widehat{v}\left(u - \sum_{i=0}^n a_i t^i\right) = \infty$$

and hence the series converges to u . Finally, we can write

$$x = ut^m = t^m(a_0 + a_1t + a_2t^2 + \dots)$$

□

We will state some results, without proof, concerning polynomials.

Let K be a complete field with respect to the valuation v . We can extend v to the ring $K[x]$ of the polynomials in one variable over K as follows:

$$v(f) := \min \{v(a_0), \dots, v(a_n)\},$$

where $f(x) = a_0 + a_1x + \dots + a_nx^n$, $a_i \in K$, $0 \leq i \leq n$, $a_n \neq 0$. A polynomial $f(x) = a_0 + a_1x + \dots + a_nx^n \in \mathcal{O}[x]$ is called **primitive** if $v(f) = 0$, i.e., $f(x) \not\equiv 0 \pmod{\mathfrak{p}}$. The following lemma is due to Kurt Hensel (1861-1941).

Lemma 7. Let $f(x) \in \mathcal{O}[x]$ be a primitive polynomial. Assume that

$$f(x) \equiv \bar{g}(x)\bar{h}(x) \pmod{\mathfrak{p}},$$

with $\bar{g}(x), \bar{h}(x) \in \mathcal{K}[x]$. Then, there exists two polynomials $g(x), h(x) \in \mathcal{O}[x]$ with $\deg g(x) = \deg \bar{g}(x)$ and

$$g(x) \equiv \bar{g}(x) \pmod{\mathfrak{p}} \text{ and } h(x) \equiv \bar{h}(x) \pmod{\mathfrak{p}},$$

such that

$$f(x) = g(x)h(x).$$

We have an immediate consequence.

Corollary. For all irreducible polynomial $f(x) = a_0 + a_1x + \cdots + a_nx^n \in K[x]$, we have

$$v(f) = \min \{v(a_0), v(a_n)\}.$$

Moreover, if $a_n = 1$ and $a_0 \in \mathcal{O}$, $f \in \mathcal{O}[x]$.

2.2 Example: the field of p -adic numbers \mathbb{Q}_p

In this example we deal, as in paragraph 1.5, with the field of rational numbers \mathbb{Q} . By theorem 5, we know that the equivalence classes of absolute values on \mathbb{Q} are represented by $|\cdot|_p$, $p > 1$ prime, and $|\cdot|$. Theorem 8 tells us that the completion of \mathbb{Q} with respect to $|\cdot|$ is isomorphic to \mathbb{R} or \mathbb{C} , since $|\cdot|$ is Archimedean. In fact, we know that this completion is \mathbb{R} (one way to define \mathbb{R} is to complete \mathbb{Q} with respect to the usual absolute value).

We are more interested in non-Archimedean absolute values. Hence, let $p > 1$ be a prime number. The completion of \mathbb{Q} with respect to $|\cdot|_p$ is denoted \mathbb{Q}_p , and called the **field of p -adic numbers**. Instead of $|\cdot|_p$, we use the corresponding valuation v_p ; we use the notation v_p also for the extension of v_p in \mathbb{Q}_p . We know that $\mathcal{K}_p \cong \mathbb{Z}/p\mathbb{Z}$, therefore we can take $\{0, \dots, p-1\}$ as set of representatives of \mathcal{K}_p ; furthermore, p is a prime element. According to theorem 10, for all $0 \neq x \in \mathbb{Q}_p$, we have

$$x = p^m(a_0 + a_1p + a_2p^2 + \cdots) = \sum_{i=m}^{\infty} a_i p^i,$$

with $a_i \in \{0, \dots, p-1\}$, $i \in \mathbb{N}$, $a_0 \neq 0$ and $m \in \mathbb{Z}$. By the construction we made in the proof of theorem 10, we know that

$$u := a_0 + a_1p + a_2p^2 + \cdots = \sum_{i=0}^{\infty} a_i p^i$$

is a unit, i.e., $v_p(u) = 0$. This means that $v_p(x) = m$. Therefore, the valuation ring of \mathbb{Q}_p is

$$\mathbb{Z}_p := \left\{ \sum_{i=m}^{\infty} a_i p^i \mid a_i \in \{0, \dots, p-1\}, a_0 \neq 0, m \geq 0 \right\},$$

called the **ring of p -adic integers**. Its group of units is

$$\mathbb{Z}_p^* = \left\{ \sum_{i=m}^{\infty} a_i T^i \mid a_i \in \{0, \dots, p-1\}, a_0 \neq 0, m = 0 \right\}$$

and the unique maximal ideal is $p\mathbb{Z}_p$. Moreover, the residue field of \mathbb{Z}_p is $\mathbb{Z}/p\mathbb{Z}$ since it is isomorphic to the residue field of \mathcal{O}_p .

Remark on notation: Note that sometimes in topology the notation \mathbb{Z}_p stands for the finite field $\mathbb{Z}/p\mathbb{Z} = \mathbb{F}_p$.

2.3 Example: the field of Laurent series $\mathbb{F}_q((T^{-1}))$

As before, we continue the example of section 1, paragraph 1.6. Let $q = p^n$, $p > 1$ prime, $n \in \mathbb{N} \setminus \{0\}$ and let $\mathbb{F}_q(T)$ be the rational function field. By theorem 6, we know that all equivalence classes of valuation on $\mathbb{F}_q(T)$ are represented by $v_{p(T)}$, $p(T) \in \mathbb{F}_q[T]$ a monic, irreducible polynomial, and v_∞ .

Now, we are going to see what happens when we complete $\mathbb{F}_q(T)$ with respect to the absolute value corresponding to v_∞ . We know that T^{-1} is a prime element for v_∞ and that the residue field \mathcal{K}_∞ of the valuation ring \mathcal{O}_∞ is isomorphic to \mathbb{F}_q , then, by theorem 10, we can write all element $f \neq 0$ of the completion in the form

$$f = (T^{-1})^m \left(a_0 + a_1 T^{-1} + a_2 (T^{-1})^2 + \dots \right) = (T^{-1})^m \sum_{i=0}^{\infty} a_i (T^{-1})^i$$

with $a_i \in \mathbb{F}_q$, $i \in \mathbb{N}$, $a_0 \neq 0$ and $m \in \mathbb{Z}$. In fact, we abuse of notations and we write

$$f = f(T) = \sum_{i=-\infty}^{-m} a_i T^i, \quad a_{-m} \neq 0.$$

We note the completion of $\mathbb{F}_q(T)$ with respect to v_∞ by $\mathbb{F}_q((T^{-1}))$. Note the analogy between $f(T) \in \mathbb{F}_q((T^{-1}))$ and a Laurent series in \mathbb{C} . Recall that a Laurent series in \mathbb{C} is a series that allows infinite negative terms and converges in an annulus. We call $\mathbb{F}_q((T^{-1}))$ the **field of formal Laurent series in T^{-1} over \mathbb{F}_q** . As above for \mathbb{Q}_p , we use the same notation for the valuation in the completion as in the rational function field. Clearly, we have $v_\infty(f(T)) = m$. The valuation ring of the field of formal Laurent series is the ring

$$\left\{ \sum_{i=-\infty}^{-m} a_i T^i \mid a_i \in \mathbb{F}_q, a_{-m} \neq 0, m \geq 0 \right\}$$

and the units are element of the form

$$\sum_{i=-\infty}^0 a_i T^i,$$

with $a_0 \neq 0$. The unique maximal ideal of the valuation ring is the set of all the series with only negative powers of T . Note that the valuation ring coincides with the ring of formal power series in T^{-1} , since no negative powers of T^{-1} occur.

Contents

1	Absolute values and valuations	1
1.1	Absolute values	1
1.2	Valuations	3
1.3	Relations between non-Archimedean absolute values and valuations .	5
1.4	Valuation ring and residue field	6
1.5	Example: the field of rational numbers \mathbb{Q}	9
1.6	Example: the rational function field $\mathbb{F}_q(T)$	12
2	Completion	17
2.1	Definitions and results	17
2.2	Example: the field of p -adic numbers \mathbb{Q}_p	21
2.3	Example: the field of Laurent series $\mathbb{F}_q((T^{-1}))$	22

References

Algebraic function fields and codes, Sections I.1 and I.2, H. Stichtenoth, Springer-Verlag, 1993.

Algebraic number theory, Sections II.3 and II.4, J. Neukirch, Springer-Verlag, 1999.